

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY
International Bureau

WO 9606078A1

INTERNATIONAL APPLICATION PUBLISHED UNDER

WO 96/06078

(51) International Patent Classification 6 :

**C07D 207/12, 207/32, 401/12, 405/12,
409/12, A61K 31/40****A1**

(11) International Publication Number:

(43) International Publication Date: 29 February 1996 (29.02.96)

(21) International Application Number:


PCT/JP94/01399

(22) International Filing Date:

24 August 1994 (24.08.94)(71) Applicant: **PFIZER PHARMACEUTICALS INC. [JP/JP];**
Shinjuku Mitsui Building, 1-1, Nishi-shinjuku 2-chome,
Shinjuku-ku, Tokyo 163-04 (JP).(72) Inventor: **ITO, Fumitaka; 1-2, Sakuragaoka 1-chome,**
Taketocho-cho, Chita-gun, Aichi 470-23 (JP).(74) Agent: **KOBAYASHI, Hiroshi; Pfizer Pharmaceuticals Inc.,**
Shinjuku Mitsui Building, 1-1, Nishi-shinjuku 2-chome,
Shinjuku-ku, Tokyo 163-04 (JP).(81) Designated State: **JP.****Published***With international search report.*(54) Title: **N-(2-(PYRROLIDINYL-1)-1-PHENYLETHYL)ACETAMIDES AS KAPPA RECEPTOR ANTAGONISTS**

(57) Abstract

A compound of formula (I) and its pharmaceutically acceptable salt, wherein R is hydrogen or hydroxy; Ar is unsubstituted or substituted phenyl; X is unsubstituted or substituted phenyl or heterocyclic, mono-, di- or trihalomethyl, cyano, or the like; and X¹ is phenyl, furyl, thienyl, pyridyl, thiazolyl, benzofuryl, benzothenyl or the like. These compounds have agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic, and neuroprotective agent.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|--|----|--------------------------|
| AT | Austria | GB | United Kingdom | MR | Mauritania |
| AU | Australia | GE | Georgia | MW | Malawi |
| BB | Barbados | GN | Guinea | NE | Niger |
| BE | Belgium | GR | Greece | NL | Netherlands |
| BF | Burkina Faso | HU | Hungary | NO | Norway |
| BG | Bulgaria | IE | Ireland | NZ | New Zealand |
| BJ | Benin | IT | Italy | PL | Poland |
| BR | Brazil | JP | Japan | PT | Portugal |
| BY | Belarus | KE | Kenya | RO | Romania |
| CA | Canada | KG | Kyrgyzstan | RU | Russian Federation |
| CF | Central African Republic | KP | Democratic People's Republic of Korea | SD | Sudan |
| CG | Congo | KR | Republic of Korea | SE | Sweden |
| CH | Switzerland | KZ | Kazakhstan | SI | Slovenia |
| CI | Côte d'Ivoire | LI | Liechtenstein | SK | Slovakia |
| CM | Cameroon | LK | Sri Lanka | SN | Senegal |
| CN | China | LU | Luxembourg | TD | Chad |
| CS | Czechoslovakia | LV | Latvia | TG | Togo |
| CZ | Czech Republic | MC | Monaco | TJ | Tajikistan |
| DE | Germany | MD | Republic of Moldova | TT | Trinidad and Tobago |
| DK | Denmark | MG | Madagascar | UA | Ukraine |
| ES | Spain | ML | Mali | US | United States of America |
| FI | Finland | MN | Mongolia | UZ | Uzbekistan |
| FR | France | | | VN | Viet Nam |
| GA | Gabon | | | | |

N-(2-(PYRROLIDINYL-1)-1-PHENYLETHYL) ACETAMIDES AS KAPPA RECEPTOR ANTAGONISTS

Technical Field

This invention relates to novel carboxamide compounds and their pharmaceutically acceptable salts, and to pharmaceutical compositions containing them. These compounds and compositions are useful as analgesic, antiinflammatory, diuretic or neuroprotective agents for the treatment of a mammalian subject, especially a human subject.


Background Art

Opioid analgesics such as morphine are therapeutically useful, but their usage is strictly limited because of their side effects such as drug dependency. Thus, analgesics with high usefulness and reduced tendency to cause drug dependency are desired. Considerable pharmacological and biochemical studies have been carried out to discover the opioid peptides and opioid receptors, and the discovery of the subtype of opioid receptor such as mu, delta, kappa at a peripheral nerve in a variety of species, including human, has made a beginning towards creating new analgesics. As it is thought that opioid analgesics such as morphine act as a μ -receptor agonist, separating the action based on a kappa-receptor agonist from the action based on μ -receptor agonist has been investigated. Recently kappa-selective agonists have been reported from the above viewpoint for example, EMD-60400: A. Barber et al., Naunyn-Schmied. Arch. Pharmacol., 345 (Suppl.): Abst 456. Some of them actually have been studied in clinical trials (Med. Res. Rev., 12, 525 (1992)).

However, even when a selective kappa-receptor agonist is employed, use of high doses can give rise to side effects such as hallucination and sedation. Therefore, it would be desired to provide compounds having better agonist activity toward opioid kappa receptor.

Brief Disclosure of the Invention

The present invention provides a compound of the following formula:



and its pharmaceutically acceptable salt, wherein


R is hydrogen or hydroxy;

Ar is phenyl or phenyl substituted with one to three substituents selected from halo, C₁₋₄ alkyl and C₁₋₄ alkoxy;

5 X is phenyl or heterocyclic; phenyl or heterocyclic substituted with one to three substituents selected from halo, C₁₋₄ alkyl, C₁₋₄ alkoxy and methoxycarbonyl; mono-, di- or tri-halomethyl; cyano; COR¹, CH=NOR², OR², SR², CH₂CN, CH₂OR², CH₂SR², CH₂S(O)R², CH₂S(O)₂R², CH₂(R²)R³, CH₂N(R²)R³, CH₂NR²OH, CH₂N(COR²)OH, CH₂NR²COR³, CH₂NR²S(O)₂R³ or
10 CH₂OCOR², wherein R¹ is hydrogen, hydroxy, amino, NHOH, NHOCH₃, pyridylamino, NHN(CH₃)₂, C₁₋₄ alkoxy, benzyloxy, C₁₋₄ alkylamino, di-C₁₋₄ alkylamino, C₁₋₄ alkyl or C₁₋₄ alkylthio; and R² and R³ are each hydrogen, C₁₋₄ alkyl, C₁₋₄ alkoxy or C₇₋₁₁ phenylalkyl; and

X¹ is phenyl, furyl, thienyl, pyridyl, thiazolyl, benzofuryl or benzothienyl;
15 phenyl, furyl, thienyl, pyridyl, thiazolyl, benzofuryl or benzothienyl, substituted with one to three substituents selected from halo, C₁₋₄ alkyl, C₁₋₄ alkoxy, amino, hydroxy, nitro, trifluoromethyl and mesyl.

Further, the present invention provides a compound of the formula:



20 wherein R, Ar and X are as already defined. These compounds can be used as intermediates to prepare the compounds of formula (I).

The carboxamide compounds of the present invention of formula (I) exhibit significant agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic and neuroprotective agents, in mammals,
25 especially man.

Accordingly, the present invention also provides a pharmaceutical

composition useful as an analgesic, antiinflammatory, diuretic or neuroprotective agent, in a mammal, especially man, which comprises a therapeutically effective amount of the carboxamide compound of formula (I) or its pharmaceutically acceptable salt together with a pharmaceutically acceptable carrier.

5

Detailed Disclosure of the Invention

In this specification, the term "heterocyclic" means a monocyclic or bicyclic hydrocarbon group which has one or more hetero atoms in the ring, preferably has 4 to 10 carbon atoms and 1 to 3 heteroatoms, including piperidino, morpholino, thiamorpholino, pyrrolidino, pyrazolino, pyrazolidino, pyrazoryl, piperaziny, furyl, 10 benzofuryl, thienyl, benzothienyl, oxazolyl, tetrazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridyl, pyrimidinyl, pyrrolyl, pyrrolidinyl, quinolyl and quinuclidinyl.

A preferred group of compounds of this invention includes the compounds of formula (I) wherein R is hydroxy; Ar is phenyl optionally substituted with one to three halogen atoms, preferably phenyl; X is phenyl optionally substituted with one 15 to three substituents selected from halo, C₁₋₄ alkyl, C₁₋₄ alkoxy and methoxycarbonyl; and X¹ is phenyl optionally substituted with one to three halogen atoms, preferably 3,4-dichlorophenyl.

Another preferred group of compounds of this invention includes the compounds of formula (I) wherein R is hydroxy; Ar is phenyl optionally substituted 20 with one to three halogen atoms, more preferably phenyl; X is mono-, di- or tri-halomethyl, cyano, hydroxycarbonyl, butyloxycarbonyl, benzyloxycarbonyl, carbamoyl or hydroxymethyl; and X¹ is phenyl optionally substituted with one to three halogen atoms, preferably 3,4-dichlorophenyl.

Another preferred group of compounds of this invention includes the 25 compounds of formula (I) wherein R is hydroxy; Ar is phenyl optionally substituted with one to three halogen atoms, more preferably phenyl; X is furyl, thienyl or pyridyl; and X¹ is phenyl optionally substituted with one to three halogen atoms, preferably 3,4-dichlorophenyl.

Preferred individual compounds of the invention are:


30 N-carboxymethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;
2-(3,4-dichlorophenyl)-N-(2-hydroxyethyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-

- phenylethyl]acetamide;
 2-(3,4-dichlorophenyl)-N-[(2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(2,2,2-trifluoroethyl)acetamide;
 2-(3,4-dichlorophenyl)-N-furfuryl-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;
 5 2-(3,4-dichlorophenyl)-N-[2-(3-(s)-hydroxypyrrolidin-1-yl)-1-(s)-phenylethyl]-N-(4-pyridyl)methylacetamide;
 2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(3-pyridyl)methylacetamide;
 10 N-cyanomethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;
 2-(3,4-dichlorophenyl)-N-(2,2-difluoroethyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;
 N-2-cyanoethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide; and
 15 2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-methoxycarbonylmethylacetamide.

General Synthesis

- The carboxamide compounds of formula (I) of this invention may be prepared
 20 by a variety of synthetic methods. For example, the carboxamide compounds of formula (I) may be prepared by acylation of compound (II), as indicated in the following Preparation Method A-I.

Preparation Method A-I:



- 25 (wherein R, Ar, X and X¹ are as previously defined)


In Preparation Method A-I, the amine compound (II) is reacted with an acylating agent using standard acylating techniques known to those skilled in the art.

In a typical acylation method, the amine compound (II) may be reacted with acyl halide (e.g., X^1CH_2COCl) in a suitable reaction-inert solvent. Suitable inert-reaction solvents include, for example, aromatic hydrocarbons such as benzene, toluene and xylene; ethers such as ethyl ether, dioxane and tetrahydrofuran; halogenated hydrocarbons such as chloroform, dichloromethane and dichloroethane; amides such as N,N-dimethylformamide; and nitriles such as acetonitrile. If desired, this reaction may be catalyzed by a base such as triethylamine, pyridine or alkoxide. The reaction may be carried out at a temperature of from $-30^\circ C$ to $100^\circ C$, preferably from $0^\circ C$ to $25^\circ C$, for 10 minutes to 48 hours, preferably from 30 minutes to 24 hours.

The compound (I) of the present invention may also be obtained from the amine compound (II) by the other acylation methods, for example, (1) a reaction with anhydride (e.g., $(X^1CH_2CO)_2O$) or a mixed anhydride in the presence of base; (2) a reaction with carboxylic acid (X^1CH_2COOH) in the presence of a coupling agent such as dicyclohexylcarbodiimide (DCC), water soluble carbodiimide (WSCD), 2-ethoxy-N-ethoxycarbonyl-1,2-dihydroquinoline, Bop agent (Benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate), diethyl azodicarboxylate-triphenylphosphine, diethyl cyanophosphonate, carbonyldiimidazole and diphenylphosphoryl azide; or (3) a reaction with carboxylic ester (e.g., X^1CH_2COOR' wherein R' is lower alkyl) optionally in the presence of base. The conditions employed for the acylation methods can be properly chosen by the skilled persons.

In an alternative method, the compound (I) of the present invention may be prepared by the following Preparation Method A-II.

Preparation Method A-II:




In this method, the compound (I) may be obtained by alkylation of the amide compound (III). Alkylation methods known to those skilled in the art can be used. For example, the amide compound (III) may be reacted with alkylhalide (e.g.,

XCH₂L wherein X is as previously defined; and L is halo such as chloro) in a reaction-inert solvent. If desired, this reaction may be catalyzed by a base such as sodium, sodium hydride, sodium hydroxide, potassium hydroxide, with or without a phase-transfer catalyst. The reaction may be carried out at a temperature of from 0°C to 200°C, preferably from 60°C to 150°C, for 5 minutes to 24 hours, preferably from 30 minutes to 12 hours.

Alternatively, the alkylation of the compound (III) may be carried out by reacting the compound (III) with formaldehyde and metal salts (e.g., MX wherein X is as previously defined; and M is an alkali metal such as sodium and potassium) in a suitable reaction-inert solvent. In addition, the amide compound (III) may be obtained by acylation of the amine compound (IV) in similar procedures to those described in Preparation Method A-I above.

In the present invention, the amine compound (II) may be obtained by the following Preparation Method B-I.

Preparation Method B-I:




In this method, the amine compound (II) may be obtained by alkylation of the amine compound (IV) using standard alkylation techniques known to those skilled in the art. A preferred alkylation method is reductive alkylation wherein the amine compound (IV) may be reacted with aldehyde, XCHO (wherein X is as already defined) in the presence of a reducing agent such as NaBH₄, NaBH₃CN or NaBH(OAc)₃. This reaction may be carried out in a suitable reaction-inert solvent at a temperature of from -20°C to 60°C, preferably from 0°C to 25°C, for 10 minutes to 48 hours, preferably from 60 minutes to 5 hours. In an alternative alkylation method, the amine compound (II) may be obtained by reacting the amine compound (IV) with alkylhalide, XCH₂L (wherein X and L are as already defined) under conditions known to those skilled in the art. The Mannich type alkylation can be also used, which comprises the reaction of the compound (IV) with formaldehyde and

a metal salt. The amine compounds (IV) are either known or may be prepared by known methods as described in European Patent No. 254545.

Alternatively, the amine compound (II) may be obtained by acylation of the compound (IV), followed by reduction, as indicated in the following Preparation

5 Method B-II.


Preparation Method B-II:



In a typical procedure, the amine compound (IV) may be first reacted with acylating agents, XCOOH (wherein X is as already defined) in the presence of a
10 suitable coupling agent as mentioned above, in a suitable reaction-inert solvent, followed by reduction using a reducing agent such as LiAlH_4 , $\text{BH}_3 \cdot \text{Me}_2\text{S}$ or $\text{BH}_3 \cdot \text{THF}$. This reaction may be carried out at a temperature of from 0°C to 100°C , preferably from 20°C to 80°C , for 30 minutes to 24 hours, preferably from 60 minutes to 12 hours. The other possible acylation methods prior to the reduction
15 include a reaction of the compound (IV) with acyl halide, XCOL in the presence of base; and the reaction of the compound (IV) with anhydride, $(\text{XCO})_2\text{O}$ in the presence of base. The conditions to be employed for these acylation methods can be appropriately chosen by those skilled in the art.

Further, the compound (II) may be obtained by acylation of an amide
20 compound of the following formula (V), followed by reduction, as indicated in the following Preparation Method B-III.

Preparation Method B-III:



In this method, the amide compound (V) may be first subjected to acylation as mentioned in the above Preparation Method B-II, and then subjected to reduction, to obtain the compound (II). The conditions for this reaction may be similar to those described in the above Preparation Method B-II. In addition, the amide compound (V) is either known or can be prepared by known methods as described in, for example, European Patent No. 254545 and Chem. Pharm. Bull., 42(3) 690-693, 1994.

The compounds of formula (I), and the intermediates shown in the above Preparation Methods can be isolated and purified by conventional procedures, such as recrystallisation or chromatographic purification.

As the carboxamide compounds of this invention possess at least two asymmetric centers, they are capable of occurring in various stereoisomeric forms or configurations. Hence, the compounds can exist in separated (+)- and (-)-optically active forms, as well as mixtures thereof. The present invention includes all such forms within its scope. Individual isomers can be obtained by known methods, such as optically selective reaction or chromatographic separation in the preparation of the final product or its intermediate.

The carboxamide compounds of the present invention can be used in the form of the inorganic salts with acid such as hydrochloric acid, hydrobromic acid, sulfonic acid, nitric acid, phosphoric acid and the like and the organic salts with acid such as acetic acid, formic acid, benzoic acid, oxalic acid, succinic acid, fumaric acid, maleic acid, citric acid, alkylsulfonic acid.

The carboxamide compounds of the present invention of formula (I) exhibit significant agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic and neuroprotective agents for the treatment of mammals, especially humans in need of such agents.

The activity of the carboxamide compounds of formula (I) of the present invention as opioid kappa agonist, is demonstrated by the opioid receptor binding activity. Such activity may be determined in homogenates from guinea pig whole brain, as described by Regina, A. et al. in J. Receptor Res. 12: 171-180, 1992. In summary, tissue homogenate is incubated at 25°C for 30 min in the presence of labelled ligand and test compounds. The m-sites are labelled by 1 nM of (3H)-[D-

Ala2,MePhe4,Gly-ol5]enkephalin (DAMGO), the d-sites by 1 nM f (3H)-[D-Pen2,5]enkephalin (DPDPE) and the k-sites by 0.5 nM (3H)-CI-977. The non specific binding is measured by use of 1 mM CI-977 (k), 1 mM (DAMGO) (m), 1mM (DPDPE) (d). Data are expressed as the IC₅₀ values obtained by a non-linear fitting
5 program using the Cheng and Prusoff equation. All compounds prepared in the Working Examples as described below were tested by this method, and showed an IC₅₀ value of 0.01 nM to 10 μ M with respect to inhibition of binding at its receptor.

The agonist activity toward opioid kappa receptor can also be demonstrated by the Formalin Test as described by Wheeler-Aceto, H. et al. in
10 Psychopharmacology 104: 35-44, 1991. In this testing, male SD rats (80-100 g) are injected s.c. with a test compound dissolved in 0.1% methyl cellulose saline or vehicle. After 30 min., 50 ml of a 2% formalin are injected into a hind paw. The number of licking the injected paw per observation period is measured 15-30 min. after the injection of formalin and expressed as % inhibition compared to the
15 respective vehicle group.

The agonist activity toward opioid kappa receptor can also be demonstrated by the Rotarod Test as described by Hayes, A.G. et al. in Br. J. Pharmacol. 79: 731-736, 1983. In this testing, a group of 6-10 male SD rats (100-120 g) are selected for their ability to balance on a rotating rod (diameter 9 cm, rate of rotation 5 r.p.m.).
20 The selected rats are then injected s.c. with a test compound dissolved in 0.1% methyl cellulose saline. The animals are tested again 30 min. after treatment; a rat falling off the bar more than twice within 150 seconds is considered to be showing motor impairment and the animal's performance (i.e., time on the rotarod) are recorded. The ED₅₀ value, defined as the dose of the drug which halves the
25 performance time is observed in the control group.

The carboxamide compounds of formula (I) of this invention can be administered via either the oral, parenteral or topical routes to mammals. In general, these compounds are most desirably administered to humans in doses ranging from 0.01 mg to 100 mg per day, although variations will necessarily occur
30 depending upon the weight and condition of the subject being treated, the disease state being treated and the particular route of administration chosen. However, a dosage level that is in the range of from 0.01 mg to 50 mg per kg of body weight per day

is most desirably employed for the treatment of pain in a postoperative patient.

The compounds of the present invention may be administered alone or in combination with pharmaceutically acceptable carriers or diluents by either of the above routes previously indicated, and such administration can be carried out in single or multiple doses. More particularly, the novel therapeutic agents of the invention can be administered in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, lozenges, troches, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various nontoxic organic solvents, etc. Moreover, oralpharmaceutical compositions can be suitably sweetened and/or flavored. In general, the therapeutically-effective compounds of this invention are present in such dosage forms at concentration levels ranging 5% to 70% by weight, preferably 10% to 50% by weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dipotassium phosphate and glycine may be employed along with various disintegrants such as starch and preferably corn, potato or tapioca starch, alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatine capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

For parenteral administration, solutions of a compound of the present invention in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably buffered (preferably pH > 8) if

necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intra-articular, intra-muscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well-known to those skilled in the art. Additionally, it is also possible to administer the compounds of the present invention topically when treating inflammatory conditions of the skin and this may preferably be done by way of creams, jellies, gels, pastes, ointments and the like, in accordance with standard pharmaceutical practice.

Examples

The present invention is illustrated by the following examples. However, it should be understood that the invention is not limited to the specific details of these examples. Melting points were taken with a Buchi micro melting point apparatus and uncorrected. Infrared Ray absorption spectra (IR) were measured by a Shimadzu infrared spectrometer (IR-470). ^1H and ^{13}C nuclear magnetic resonance spectra (NMR) were measured in CDCl_3 by a JEOL NMR spectrometer (JNM-GX270, 270MHz) unless otherwise indicated and peak positions are expressed in parts per million (ppm) downfield from tetramethylsilane. The peak shapes are denoted as follows: s, singlet; d, doublet; t, triplet; m, multiplet; br, broad.

Preparation 1

(S)-Phenylglycyl-3-(S)-hydroxypyrrolidine

To a stirred solution of 3-(S)-pyrrolidinol (3.054g, 35mmol) and N-benzyloxycarbonyl-(S)-phenylglycine (10.00g, 35mmol) in DMF (40ml) was added diethyl phosphorocyanidate (6.28ml, 42mmol) followed by addition of N-methylmorpholine (4.65ml, 42mmol) at room temperature. After 1h stirring at room temperature, the reaction mixture was poured into water (200ml), extracted with mixed solvent (ethyl acetate / hexane / ether : 2/1/1, 100ml x 3). The extract combined was washed with 1N HCl solution, saturated NaHCO_3 aqueous solution and brine, dried (MgSO_4), and concentrated to give 11.395g (91.9%) of N-benzyloxycarbonyl-(S)-phenylglycyl-3-(S)-hydroxypyrrolidine as yellow oil. A suspension mixture of this oil (11.395g, 32mmol) and 10% palladium carbon (1.14g) in methanol (100ml) was stirred under hydrogen atmosphere at room temperature for

17h. The catalyst was removed by Celite filtration and the filtrate was concentrated to give 8.255g (crude 100%) of the title compound as brown oil.

^1H NMR (270MHz, CDCl_3) δ 7.45-7.27 (5H, m), 4.56 (0.7H, s), 4.49 (0.3H, s), 4.45-4.35 (1H, m), 3.74-3.36 (3H, m), 3.25-3.15 (0.7H, m), 3.10-2.98 (0.3H, m),
5 2.70 (2H, br.s), 2.30 (1H, br.s), 2.05-1.75 (2H, m).
IR(neat) : 3350, 3300, 1640 cm^{-1} .

To a stirred solution of 3-(S)-pyrrolidinol (14.70g, 169mmol) and N-t-butoxycarbonyl-(S)-phenylglycine (42.46g, 169mmol) in DMF (250ml) was added diethyl phosphorocyanidate (30.3ml, 203mmol) followed by addition of N-methylmorpholine (22.5ml, 203mmol) at room temperature. After 1h stirring at
10 room temperature, the reaction mixture was poured into water (1250ml), extracted with mixed solvent (ethyl acetate/hexane/ether:2/1/1, 400ml x 3). The extract combined was washed with 1N HCl solution, saturated NaHCO_3 aqueous solution and brine, dried (MgSO_4), and concentrated to give 34.81g (64.3%) of N-t-
15 butoxycarbonyl-(S)-phenylglycyl-3-(S)-hydroxypyrrolidine as white powder. To a stirred suspension of N-t-butoxycarbonyl-(S)-phenylglycyl-3-(S)-hydroxypyrrolidine (10.13g, 31.6mmol) in CH_2Cl_2 (10ml) was added trifluoroacetic acid (25ml) at 0 and resulting solution was stirred at room temperature for 1h. After evaporation of excess trifluoroacetic acid and solvent, the residue was basified with aqueous NH_3 solution
20 (20ml) and extracted with CH_2Cl_2 (30ml x 3). After dry (Na_2SO_4), the solvent was evaporated to give 4.974g (71.5%) of title compound.

Preparation 2

(2S,3S)-1-(2-Amino-2-phenylethyl)-3-hydroxypyrrolidine

To a stirred suspension of lithium aluminum hydride (3.795g, 100mmol) in
25 THF (200 ml) was added a solution of (S)-phenylglycyl-3-(S)-hydroxypyrrolidine (7.049g, 32mmol) in THF (100ml) dropwise at room temperature. The reaction mixture was refluxed for 1.5h. Then the reaction mixture was cooled down to room temperature and $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ (10.31g) and KF (1.86g) was added to the reaction mixture. After 1h stirring, the white solid precipitated was removed by Celite
30 filtration and the filtrate was concentrated to give 4.21g of clear yellow oil. This was purified by column chromatography (silica gel: 180g, $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$: 50/5/1 as eluent) to afford 3.584g (54.3%) of clear yellow oil.

^1H NMR (270MHz, CDCl_3) δ 7.38-7.15 (5H, m), 4.37-4.29 (1H, m), 4.08 (1H, dd, $J = 4.0, 10.3\text{Hz}$), 3.08-3.01 (1H, m), 2.77 (1H, dd, $J = 10.3, 12.1\text{Hz}$), 2.69-2.61 (1H, m), 2.43 (1H, dd, $J = 4.0, 12.1\text{Hz}$), 2.31-2.00 (6H, m), 1.83-1.70 (1H, m).
IR (neat) : 3350, 3200 cm^{-1} .

5

Preparation 3

(2S,3S)-1-[2-N-(Benzyloxycarbonyl)methylamino-2-phenylethyl]-3-hydroxypyrrolidine

A mixture of (2S,3S)-1-(2-amino-2-phenylethyl)-3-hydroxypyrrolidine (1.65g, 8mmol), benzyl 2-bromoacetate (1.52ml, 9.6mmol), and triethylamine (1.34ml, 9.6mmol) in CH_2Cl_2 (30ml) was refluxed for 7h. The reaction mixture was diluted with water (50ml), then extracted with CH_2Cl_2 (50ml x 3). The extract combined was washed with brine and dried (MgSO_4). Evaporation of the solvent gave 2.18g of brown oil which was purified by column chromatography (silica gel: 70g, $\text{CH}_2\text{Cl}_2/\text{MeOH}$: 20/1 to 10/1 as eluent) to afford 1.218g (42.9%) of clear yellow oil.

15

^1H NMR (270MHz, CDCl_3) δ 7.40-7.22 (10H, m), 5.13 (2H, s), 4.35-4.25 (1H, m), 3.82 (1H, dd, $J = 3.7, 11.0\text{Hz}$), 3.43 (1H, d, $J = 17.6\text{Hz}$), 3.22 (1H, d, $J = 17.6\text{Hz}$), 3.16-3.06 (1H, m), 2.86 (1H, dd, $J = 11.0, 12.1\text{Hz}$), 2.70 (1H, dd, $J = 4.8, 10.3\text{Hz}$), 2.64 (1H, dd, $J = 1.8, 9.9\text{Hz}$), 2.31 (1H, dd, $J = 3.7, 12.1\text{Hz}$), 2.26-2.15 (4H, m), 1.82-1.71 (1H, m).
IR (neat) : 3350, 1740 cm^{-1} .

20

Example 1

N-(Benzyloxycarbonyl)methyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

25

To a stirred solution of (2S,3S)-1-[2-N-(benzyloxycarbonyl)methylamino-2-phenylethyl]-3-hydroxypyrrolidine (354g, 1mmol) in dioxane (4ml) was added 1N NaOH aqueous solution (1 ml) followed by dropwise addition of 3, 4-dichlorophenylacetyl chloride at room temperature. After 2.5h stirring, the reaction mixture was extracted with CH_2Cl_2 . The extract combined was washed with brine, dried (MgSO_4), and concentrated to give 0.952g of brown viscous oil, which was purified by column chromatography (silica gel : 30g, $\text{CH}_2\text{Cl}_2/\text{MeOH}$: 30/1 to 10/1 as eluent) to afford 152mg (28.1%) of title compound as yellow brown viscous oil.

30

¹H NMR (270MHz, CDCl₃) δ 7.39-7.27 (11H, m), 7.15-7.08 (1.5H, m), 7.01 (0.5H, dd, J = 2.2, 8.4Hz), 6.04 (0.5H, dd, J = 5.9, 9.9Hz), 5.10 (1H, s), 5.07 (0.5H, d, J = 12.1Hz), 5.00 (0.5H, t, J = 7.5Hz), 4.99 (0.5H, d, J = 12.1Hz), 4.28 (0.5H, d, J = 16.8Hz), 4.30-4.20 (1H, m), 3.97 (0.5H, d, J = 18.7Hz), 3.92 (0.5H, d, J = 16.5Hz), 3.89 (0.5H, d, J = 15.7Hz), 3.76 (1H, s), 3.71 (0.5H, d, J = 16.1Hz), 3.60 (0.5H, d, J = 15.7Hz), 3.14 - 2.42 (5H, m), 2.27 - 2.01 (2H, m), 1.80 - 1.60 (2H, m).
 IR (neat) : 3450, 1750, 1650 cm⁻¹.

This oil was converted to HCl salt using HCl gas saturated methanol to give 77mg of amorphous solid.
 Anal. Calcd for C₂₉H₃₀Cl₂N₂O₄•HCl•1.5H₂O : C, 57.58 ; 5.66; N, 4.63 Cl, 17.58.
 Found : C, 57.56 ; H, 5.37 ; N, 4.71 ; Cl, 17.67.

Example 2

(2S,3S)-1-[2-N-(t-Butoxycarbonyl) methylamino-2-phenylethyl]-3-hydroxypyrrolidine

This compound was prepared in 70% yield according to a procedure similar to that described in Preparation 3.

¹H NMR (270MHz, CDCl₃) δ 7.34-7.28 (5H, m), 4.38-4.30 (1H, m), 3.81 (1H, dd, J = 3.7, 10.6Hz), 3.25 (1H, d, J = 17.6Hz), 3.19-3.13 (1H, m), 3.04 (1H, d, J = 17.6Hz), 2.87 (1H, dd, J = 10.6, 12.1Hz), 2.75-2.65 (2H, m), 2.34 (1H, dd, J = 3.7, 11.7Hz), 2.27-2.15 (4H, m), 1.85-1.75 (1H, m), 1.44 (9H, s).
 IR(neat) : 3300, 1730cm⁻¹.

Example 3

N-(t-Butoxycarbonyl)methyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This compound was prepared in 37.1% yield according to a procedure similar to that described in Example 1.

¹H NMR (270MHz, CDCl₃) δ 7.41-7.10 (8H, m), 6.10 (0.7H, dd, J = 6.6, 9.6Hz), 5.03 (0.3H, dd, J = 7.0, 7.7Hz), 4.30-4.20 (1H, m), 3.93 (0.3H, d, J = 16.8Hz), 3.82 (0.6H, s), 3.82 (0.7H, d, J = 18.7Hz), 3.76 (0.3H, d, J = 15.4Hz), 3.71 (0.7H, d, J = 18.7Hz), 3.62 (0.7H, d, J = 15.8Hz), 3.55 (0.7H, d, J = 15.8Hz), 3.17-2.05 (9H, m), 1.80-1.65 (1H, m), 1.39 (2.7H, s), 1.30 (6.3H, s).

IR(neat) : 3400, 1740, 1650cm⁻¹.

Anal. Calcd for C₂₆H₃₂Cl₂N₂O₄•0.5H₂O : C, 60.47 ; H, 6.44 ; N, 5.42.

Found : C, 60.24 ; H, 6.41 ; N, 5.24.

Example 4

5 N-Carboxymethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

A mixture of N-(t-butoxycarbonyl)methyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide (306mg, 0.6mmol), HCl gas saturated methanol solution (8ml), and methanol (2ml) was refluxed with stirring for
10 2h. After evaporation of the solvent, the resulting oil was crystallized from CH₂Cl₂/MeOH to give 153mg (52.3%) of desired compound as HCl salt of white powder.

mp 161.9 - 163.5

Anal. Calcd for C₂₂H₂₄Cl₂N₂O₄•HCl : C, 54.17 ; H, 5.17 ; N, 5.74.

15 Found : C, 54.40 ; H, 5.47 ; N, 5.62.

¹H NMR (270MHz, CDCl₃) δ 11.30 (1H, br.s), 7.45-7.25 (8H, m), 6.45 (1H, m), 4.65-2.00(15H, m).

IR (KBr) : 3300, 1740, 1665cm⁻¹.

Example 5

20 (2S,3S)-1-[2-N-(2-Hydroxyethylamino)-2-phenylethyl]-3-hydroxypyrrolidine

To a stirred suspension of lithium aluminum hydride (380mg, 10mmol) in THF (10ml) was added a solution of (2S, 3S)-1-[2-N-(t-butoxycarbonyl)methylamino-2-phenylethyl]-3-hydroxypyrrolidine (1.602g, 5mmol) in THF (15ml) dropwise at room temperature. The reaction mixture was then refluxed with stirring for 1h.
25 After cooling down to room temperature, Na₂SO₄•10H₂O(3.80g) and KF(0.38g) was added to the reaction mixture. After 1h stirring, the solid appeared was removed by Celite filtration. The filtrate was concentrated to give 1.32g(crude 100%) of desired compound as yellow oil.

¹H NMR (270MHz, CDCl₃) δ 7.34-7.24 (5H, m), 4.33 (1H, br.s), 3.77 (1H, dd, J = 3.7, 11.0Hz), 3.77-3.57 (2H, m), 3.20-3.05 (1H, m), 2.88 (1H, dd, J = 11.0, 12.1Hz), 3.05-2.55 (6H, m), 2.35 (1H, dd, J = 3.7, 12.1Hz), 2.30-2.15 (2H, m), 1.95-1.75 (2H, m).

30

1R (neat) : 3350cm⁻¹.

Example 6

2-(3,4-Dichlorophenyl)-N-(2-hydroxyethyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

5 This compound was prepared in 48% yield according to a procedure similar to that described in Example 1.

¹H NMR (270MHz, CDCl₃) δ 7.38-7.02 (8H, m), 6.13 (0.75H, br.d, J = 9.2Hz), 5.05 (0.25H, m), 4.45-4.35 (1H, m), 4.10-3.10 (9H, m), 2.90 -2.15 (6H, m), 1.90-1.75 (1H, m).

10 1R (neat) : 3400, 1640 cm⁻¹.

HCl salt : amorphous solid.

Anal. Calcd for C₂₂H₂₆Cl₂N₂O₃•HCl•2.5H₂O : C, 50.93 ; H, 6.22 ; N, 5.40.
Found : C, 51.20 ; H, 6.02 ; N, 5.66.

Example 7

15 (2S,3S)-3-Hydroxy-1-[2-phenyl-2-N-(2,2,2-trifluoroethylamino)ethyl]pyrrolidine

To a stirred solution of (S)-phenylglycyl-3-(S)-hydroxypyrrolidine (1.00g, 4.5mmol) in THF (20ml) was added trifluoroacetic anhydride (0.7ml, 5mmol) at room temperature. After 1h stirring, the solvent was evaporated and the residue was purified by column chromatography (silica gel:100g, CH₂Cl₂/MeOH:20/1 as eluent)
20 to give 0.90g of amide derivative, which was dissolved in THF (14ml) followed by addition of boran-methyl sulfide complex (1.33ml, 14mmol) at 0. Then the reaction mixture was refluxed for 13h. To this reaction mixture was added 1N HCl aqueous solution (10ml) at 0 and the mixture was refluxed for 1h. After cooling down to room temperature, the reaction mixture was basified with 1N NaOH aqueous solution
25 and extracted with CH₂Cl₂. After dry (Na₂SO₄), the solvent was evaporated to give 0.821g of colorless oil which was soon crystallized.

¹H NMR (270MHz, CDCl₃) δ 7.38-7.20 (5H, m), 4.38-4.30 (1H, m), 3.92 (1H, dd, J = 3.3, 11.0Hz), 3.13-2.95 (3H, m), 2.81 (1H, dd, J = 11.0, 12.1Hz), 2.85-2.60 (3H, m), 2.38-2.11 (3H, m), 1.95 (1H, br.s), 1.85-1.65 (1H, m).

30 1R(neat) : 3350cm⁻¹.

Example 8

2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-

(2,2,2-trifluoroethyl)acetamide

This compound was prepared in 38.9% yield according to a procedure similar to that described in Example 1.

¹H NMR (270MHz, CDCl₃) δ 7.40 (1H, d, J = 8.1Hz), 7.36-7.26 (5H, m), 7.16 (1H, br.s), 7.07 (1H, br.d, J = 8.1Hz), 5.78 and 5.06 (total 1H, each br.s), 4.27 (1.5H, br.s), 3.90-3.60 (3.5H, m), 3.80-2.80 (3H, m), 2.75-2.60 (2H, m), 2.45-2.05 (2H, m), 1.95-1.60 (2H, m).

IR (neat) : 3450, 1660 cm⁻¹.

HCl salt : amorphous solid

Anal. Calcd for C₂₂H₂₃Cl₂F₃N₂O₂•HCl : C, 51.63 ; H, 4.73 ; N, 5.47.

Found : C, 51.78 ; H, 5.19 ; N, 5.28.

Example 9(2S,3S)-1-[2-N-(2-furyl)methylamino-2-phenylethyl]-3-hydroxypyrrolidine

A mixture of (2S,3S)-1-(2-amino-2-phenylethyl)-3-hydroxypyrrolidine (619mg, 3mmol), and 2-furaldehyde (0.37ml, 4.5mmol) in ethanol (8ml) was refluxed with stirring for 4.5h. Then the solvent was evaporated and the resulting Schiff base was dissolved in methanol (10ml) and to this solution was added NaBH₄ by portions at room temperature. After 0.5h stirring at room temperature, the solvent was evaporated. Then water (30ml) was added to the residue and extracted with CH₂Cl₂. The extract was washed with brine, dried (MgSO₄), and concentrated to give 857mg (99.8%) of brown oil.

¹H NMR (270MHz, CDCl₃) δ 7.40-7.28 (6H, m), 6.30 (1H, dd, J = 1.8, 2.9Hz), 6.08 (1H, d, J = 2.9Hz), 4.33-4.25 (1H, m), 3.75 (1H, d, J = 14.7Hz), 3.71 (1H, dd, J = 3.7, 11.4Hz), 3.49 (1H, d, J = 14.7Hz), 2.84 (1H, dd, J = 11.4, 12.1Hz), 2.83-2.75 (1H, m), 2.62-2.51 (2H, m), 2.35 (2H, br.s), 2.27 (1H, dd, J = 3.7, 12.1Hz), 2.21-2.10 (2H, m), 1.80-1.65 (1H, m).

IR (neat) : 3300cm⁻¹.

Example 102-(3,4-Dichlorophenyl)-N-furfuryl-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This compound was prepared in 96.5% yield according to a procedure similar to that described in Example 1.

- ¹H NMR (270MHz, CDCl₃) δ 7.37 (1H, d, J = 8.1Hz), 7.37 (1H, d, J = 1.8Hz), 7.31-7.22 (5H, m), 7.10 (1H, dd, J = 1.8, 8.1Hz), 7.03 (1H, dd, J = 1.8, 8.1Hz), 6.19 (1H, dd, J = 1.9, 2.9Hz), 5.94 (1H, dd, J = 5.1, 9.5Hz), 5.71 (1H, d, J = 2.9Hz), 4.40-4.33 (1H, m), 4.30 (1H, d, J = 17.9Hz), 4.22 (1H, d, J = 17.9Hz), 3.72 (1H, d, J = 13.2Hz), 3.66 (1H, d, J = 13.2Hz), 3.59 (1H, dd, J = 9.5, 12.8Hz), 3.49 (1H, s), 3.45-3.35 (1H, m), 3.20-3.05 (2H, m), 2.83 (1H, dd, J = 5.1, 11.0Hz), 2.65-2.55 (1H, m), 2.55-2.10 (1H, m), 1.95-1.80 (1H, m).
1R (neat) : 3400, 1650 cm⁻¹.
HCl salt : mp 181.0 - 183.5
- 10 Anal. Calcd for C₂₅H₂₆Cl₂N₂O₃ : C, 58.89 ; H, 5.34 ; N, 5.49.
Found : C, 58.58 ; H, 5.61 ; N, 5.63.

Example 11

(2S,3S)-1-[2-Phenylethyl]-2-N-(2-thienyl)methylamino]-3-hydroxypyrrolidine

- This compound was prepared in 100% yield according to a procedure similar to that described in Example 9.
- 15 ¹H NMR (270MHz, CDCl₃) δ 7.42-7.24 (5H, m), 7.20 (1H, dd, J = 1.1, 5.1Hz), 6.94 (1H, dd, J = 3.3, 5.1Hz), 6.83 (1H, br.d, J = 2.9Hz), 4.30-4.20 (1H, m), 3.90 (1H, d, J = 14.3Hz), 3.79 (1H, dd, J = 3.7, 11.0Hz), 3.73 (1H, d, J = 14.3Hz), 2.84 (1H, dd, J = 11.0, 12.1Hz), 2.88-2.77 (1H, m), 2.62-2.52 (2H, m), 2.29 (1H, dd, J = 3.7, 12.1Hz), 2.32-2.10 (4H, m), 1.77-1.64 (1H, m).
20 1R (neat) : 3300cm⁻¹.

Example 12

2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(2-thienyl)methylacetamide

- 25 This compound was prepared in 62.6% yield according to a procedure similar to that described in Example 1.
- ¹H NMR (270MHz, CDCl₃) δ 7.39 (1H, d, J = 2.2Hz), 7.33 (1H, d, J = 8.8Hz), 7.30-7.26 (3H, m), 7.15-7.09 (3H, m), 6.94 (1H, dd, J = 2.2, 8.4Hz), 6.83 (1H, dd, J = 3.7, 5.1Hz), 6.61 (1H, br.d, J = 2.9Hz), 5.99 (1H, dd, J = 5.1, 9.5Hz), 4.59 (1H, d, J = 17.9Hz), 4.47 (1H, d, J = 17.6Hz), 4.43-4.35 (1H, m), 3.67-3.35 (5H, m), 3.21-3.08 (2H, m), 2.87 (1H, dd, J = 5.1, 11.0Hz), 2.68-2.56 (1H, m), 2.26-2.14 (1H, m), 1.96-1.85 (1H, m).
- 30

1R (neat) : 3400, 1650cm⁻¹.

HCl salt : mp 180.7 - 184.0

Anal. Calcd for C₂₅H₂₆Cl₂N₂O₂S HCl•0.4H₂O : C, 56.32 ; H, 5.26 ; N, 5.25.

Found : C, 56.67 ; H, 5.38 ; N, 4.77.

5

Example 13

(2S,3S)-1-[2-Phenylethyl-2-N-(4-pyridyl)methylamino]-3-hydroxypyrrolidine

This compound was prepared in 63.9% yield according to a procedure similar to that described in Example 9.

¹H NMR (270MHz, CDCl₃) δ 8.49(1H, s), 8.48 (1H, d, J = 4.8Hz), 7.63 (1H, dd, J = 1.5, 7.7Hz), 7.42-7.27 (5H, m), 7.24 (1H, dd, J = 4.8, 7.7Hz), 4.35-4.25 (1H, m), 3.74 (1H, dd, J = 3.7, 11.0Hz), 3.73 (1H, d, J = 13.8Hz), 3.55 (1H, d, J = 13.6Hz), 2.85 (1H, dd, J = 11.0, 12.1Hz), 2.88-2.82 (1H, m), 2.60 (2H, d, J = 4.0Hz), 2.42 (2H, br.s), 2.31 (1H, dd, J = 3.7, 12.1Hz), 2.27-2.09 (2H, m), 1.80-1.65 (1H, m).

15 1R (neat) : 3300cm⁻¹.

Example 14

2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(3-pyridyl)methylacetamide

20 This compound was prepared in 56.7% yield according to a procedure similar to that described in Example 1.

¹H NMR (270MHz, CDCl₃) δ 8.51 (0.7H, s), 8.48 (0.7H, d, J = 4.4Hz), 8.40 (0.3H, d, J = 2.9Hz), 8.26 (0.3H, s), 7.55-6.99 (10H, m), 6.26 (0.7H, dd, J = 4.0, 11.0Hz), 5.17 (0.3H, t), 4.70 (0.3H, d), 4.41-4.20 (2.7H, m), 3.90 (0.6H, s), 3.58 (0.7H, d), 3.50(0.7H, d), 3.25-2.35 (5H, m), 2.28-1.95 (2H, m), 1.85-1.60 (2H, m).

25 1R (neat) : 3300, 1650 cm⁻¹.

Anal. Calcd for C₂₆H₂₇Cl₂N₃O₂•2HCl•1.5H₂O: C, 53.44 ; H, 5.52 ; N, 7.19.

Found : C, 53.17 ; H, 5.21 ; N, 6.91.

Example 15

(2S,3S)-1-[2-Phenylethyl-2-N-(2-pyridyl)methylamino]-3-hydroxypyrrolidine

30 This compound was prepared in 37.9% yield according to a procedure similar to that described in Example 9.

¹H NMR (270MHz, CDCl₃) δ 8.56 (1H, br.d, J = 4.8Hz), 7.61 (1H, dt, J = 1.8,

7.7Hz), 7.42-7.13 (7H, m), 4.33-4.25 (1H, m), 3.83 (1H, d, $J = 14.3\text{Hz}$), 3.73 (1H, dd, $J = 3.7, 10.6\text{Hz}$), 3.67 (1H, d, $J = 14.3\text{Hz}$), 3.00- 2.80 (3H, m), 2.92 (1H, dd, $J = 10.6, 12.1\text{Hz}$), 2.70 (1H, br.d, $J = 9.5\text{Hz}$), 2.59 (1H, dd, $J = 4.4, 9.9\text{Hz}$), 2.36 (1H, dd, $J = 3.7, 12.1\text{Hz}$), 2.25-2.09 (2H, m), 1.85-1.70 (1H, m).

5 1R (neat) : 3300cm^{-1} .

Example 16

2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(2-pyridyl)methylacetamide

This compound was prepared in 68.5% yield according to a procedure similar to that described in Example 1.

^1H NMR (270MHz, CDCl_3) δ 8.79 (1H, br.s), 7.66-7.60 (1H, m), 7.33 (1H, d, $J = 8.4\text{Hz}$), 7.33-7.23 (6H, m), 7.18 (1H, d, $J = 1.8\text{Hz}$), 7.05 (1H, br.d, $J = 7.7\text{Hz}$), 6.97 (1H, dd, $J = 1.8\text{Hz}, 8.1\text{Hz}$), 6.50 (1H, dd, $J = 3.7, 12.1\text{Hz}$), 4.70-4.55 (2H, m), 4.37 (1H, d, $J = 18.3\text{Hz}$), 4.20 (1H, dd, $J = 12.5, 12.8\text{Hz}$), 3.95-3.25 (8H, m), 2.50-2.30 (1H, m), 2.25-2.10 (1H, m).

1R (neat) : $3400, 1650\text{cm}^{-1}$.

HCl Salt : mp 126.5 - 132.0

Anal. Calcd for $\text{C}_{26}\text{H}_{27}\text{Cl}_2\text{N}_3\text{O}_2 \cdot 2\text{HCl} \cdot 2\text{H}_2\text{O}$: C, 52.63 ; H, 5.61 ; N, 7.08

Found : C, 52.31 ; H, 5.39 ; N, 6.75.

20

Example 17

(2S,3S)-1-[2-Phenylethyl-2-N-(4-pyridyl)methylaminol-3-hydroxypyrrolidine

This compound was prepared in 81.8% yield according to a procedure similar to that described in Example 9.

^1H NMR (270MHz, CDCl_3) δ 8.52 (2H, d, $J = 5.5\text{Hz}$), 7.40-7.25 (5H, m), 7.22 (2H, d, $J = 5.5\text{Hz}$), 4.37-4.27 (1H, m), 3.72 (1H, d, $J = 14.3\text{Hz}$), 3.75-3.70 (1H, m), 3.56 (1H, d, $J = 14.7\text{Hz}$), 2.86 (1H, t, $J = 11.4\text{Hz}$), 2.90-2.82 (1H, m), 2.62 (2H, app d), 2.39 (2H, br.s), 2.35-2.10 (3H, m), 1.85-1.68 (1H, m).

1R (neat) : 3300cm^{-1} .

Example 18

2-(3,4-Dichlorophenyl)-N-[2-(3-(s)-hydroxypyrrolidin-1-yl)-1-(s)-phenylethyl]-N-(4-pyridyl)methylacetamide

This compound was prepared in 10.9% yield according to a procedure similar

to that described in Example 1.

¹H NMR (270MHz, CDCl₃) δ 8.48 (1.4H, d, J = 5.9Hz), 8.40 (0.6H, d, J = 5.9Hz), 7.44-6.97 (10H, m), 6.26 (0.7H, dd, J = 4.4, 11.0Hz), 5.17 (0.3H, app t), 4.62 (0.3H, d, J = 16.1Hz), 4.40-4.30 (1H, m), 4.29 (1.4H, s), 4.26 (0.3H, d, J = 16.9Hz), 3.92 (0.6H, s), 3.49 (0.7H, d, J = 15.4Hz), 3.41 (0.7H, d, J = 15.4Hz), 3.20-2.45 (5H, m), 2.35-2.00 (2H, m), 1.92 (1H, br.s), 1.85-1.65 (1H, m).
IR (neat) : 3400, 1650cm⁻¹.

HCl salt : mp 161.1 - 164.2

Anal. Calcd for C₂₆H₂₇Cl₂N₃O₂•2HCl•3H₂O : C, 51.08 ; H, 5.77 ; N, 6.87.

Found : C, 50.75 ; H, 5.26 ; N, 6.83.

Example 19

(2S,3S)-1-[2-N-(4-Methoxycarbonylphenyl)methylamino-2-phenylethyl]-3-hydroxypyrrolidine

This compound was prepared in 81.3% yield according to a procedure similar to that described in Example 9.

¹H NMR (270MHz, CDCl₃) δ 7.99 (2H, d, J = 8.1Hz), 7.40-7.29 (7H, m), 4.35-4.25 (1H, m), 3.91 (3H, s), 3.79 (1H, d, J = 13.9Hz), 3.70 (1H, dd, J = 3.3, 10.6Hz), 3.57 (1H, d, J = 13.9Hz), 2.85 (1H, dd, J = 10.6, 12.1Hz), 2.85-2.81 (1H, m), 2.62-2.52 (2H, m), 2.30 (1H, dd, J = 3.3, 12.1Hz), 2.27-2.00 (4H, m), 1.80-1.65 (1H, m).

IR (neat) : 3280, 3200, 1720cm⁻¹.

Example 20

2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(4-methoxycarbonylphenyl)-methylacetamide

This compound was prepared in 78.5% yield according to a procedure similar to that described in Example 1.

¹H NMR (270MHz, CDCl₃) δ 7.83 (2H, d, J = 8.1Hz), 7.39-6.90 (10H, m), 6.30-6.15 (1H, m), 4.60-4.10 (3H, m), 3.90 (3H, s), 3.68 (1H, t, J = 11.7Hz), 3.63-3.45 (2H, m), 3.30-3.17 (2H, m), 3.10-2.95 (2H, m), 2.74 (1H, br.s), 2.35-2.20 (1H, m), 2.05-1.90 (1H, m).

IR (neat) : 3400, 1720, 1650cm⁻¹.

HCl salt : amorphous solid

Anal. Calcd for $C_{29}H_{31}Cl_2N_2O_4 \cdot HCl \cdot 0.7H_2O$: C, 58.88 ; H, 5.69 ; N, 4.74.

Found : C, 58.56 ; H, 5.24 ; N, 4.65.

Example 21

(2S,3S)-1-(2-N-Cyanomethylamino-2-phenylethyl)-3-hydroxypyrrolidine

5 To a stirred solution of sodium bisulfite (1.249g, 12mmol) in water (1ml) was added 37% aqueous solution of formaldehyde (0.9ml, 12mmol) at 0 and the mixture was stirred at room temperature for 0.5h. Then to this mixture was added (2S,3S)-1-(2-phenylethyl)-3-hydroxypyrrolidine (2.48g, 12mmol) and the resulting mixture was stirred at 50 for 0.5h. After cooling down to room temperature, aqueous solution of
10 KCN (781.4mg, 12mmol) was added to the reaction mixture and stirring was continued for 2h. The reaction mixture was diluted with water (30ml), extracted with CH_2Cl_2 (20ml x 3). The extract was washed with saturated $NaHCO_3$ aqueous solution and dried (Na_2SO_4). The solvent was evaporated to give 2.20g of yellow oil, which was purified by column chromatography (silica gel ; 70g, CH_2Cl_2 /MeOH
15 : 50/1 as eluent) to afford 1.725g (58.6%) of colorless viscous oil which was gradually crystallized.

1H NMR (270MHz, $CDCl_3$) δ 7.39-7.27 (5H, m), 4.40-4.35 (1H, m), 3.96 (1H, dd, $J = 3.3, 11.4$ Hz), 3.67 (1H, d, $J = 17.6$ Hz), 3.24 (1H, d, $J = 17.6$ Hz), 3.17-3.09 (1H, m), 2.87 (1H, app t, $J = 11.7$ Hz), 2.75 (1H, dd, $J = 4.8, 9.9$ Hz), 2.67 (1H,
20 br.d, $J = 8.4$ Hz), 2.36 (1H, dd, $J = 3.3, 12.1$ Hz), 2.31-2.10 (3H, m), 1.90 (1H, br.s), 1.88 -1.75 (1H, m).

IR (neat) : 3300, 2230 cm^{-1} .

Example 22

N-Cyanomethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

25 This compound was prepared in 46.6% yield according to a procedure similar to that described in Example 1.

1H NMR (270MHz, $CDCl_3$) δ 7.38-7.00 (8H, m), 6.09 (0.7H, br.s), 5.31 (0.3H, br.s), 5.09 (1H, br.s), 4.45 (1H, Br.d, $J = 16.9$ Hz), 4.40-4.25 (1H, m), 4.15-3.70
30 (4H, m, including 1H, br.d, $J = 16.1$ Hz at 3.95ppm), 3.65-2.55 (4H, m), 2.50-1.70 (4H, m).

^{13}C NMR ($CDCl_3$) δ 171.2, 137.4, 134.0, 132.7, 131.4, 131.1, 130.6,

129.3, 128.6, 127.3, 126.3, 116.7, 71.2, 63.2, 59.9, 57.9, 51.9, 39.4, 35.0, 31.0.

IR (neat) : 3450, 2250, 1660cm⁻¹.

MS m/e (%) : 432 (<4)[M+], 189 (22), 161 (93), 159 (100), 145 (39), 143 (35), 132 (32), 125 (31), 123 (37), 117 (30), 100 (99).

5

Preparation 4

3-(S)-Methoxymethoxypyrrolidine

To a stirred solution of N-benzyl-3-(S)-hydroxypyrrolidine (4.785g, 27mmol) in THF (50ml) was added sodium hydride (60% oil suspension, 1.12g, 28mmol) by portions under nitrogen at room temperature. The suspension mixture was then refluxed for 1h. To this reaction mixture was added chloromethyl methyl ether (2.3ml, 30mmol) and refluxing was continued for 13h. After cooling down to room temperature, water (10ml) was added to the reaction mixture to give a solution mixture, which was basified with 1N NaOH aqueous solution, extracted with ethyl acetate (30ml x 3), and dried (Na₂SO₄). Evaporation of the solvent gave 6.33g of brown oil, which was purified by column chromatography (silica gel : 150g, CH₂Cl₂/MeOH : 20/1 as eluent) to afford 5.09g (85%) of clear brown oil. A suspension mixture of this oil (5.09g, 23mmol) and Pearlman's catalyst (2.00g) in MeOH (100ml) was stirred under hydrogen atmosphere at room temperature for 15h. After removal of the catalyst by Celite filtration, the filtrate was concentrated to give an oil and water mixture, which was dried in vacuo to afford 2.76g (91.4%) of orange color oil.

15

¹H NMR (270MHz, CDCl₃) δ 4.65 (1H, d J = 7.0Hz), 4.63 (1H, d, J = 6.6Hz), 4.30-4.24 (1H, m), 3.37 (3H, s), 3.21 (1H, br.s), 3.17-2.86 (4H, m), 1.98-1.80 (2H,m).

25

IR(neat) : 3300cm⁻¹.

Preparation 5

(S)-Phenylglycyl-3-(S)-methoxymethoxypyrrolidine

This compound was prepared in 61.8% yield using 3-(S)-methoxymethoxypyrrolidine and N-benzyloxycarbonyl-(S)-phenylglycine according to a procedure similar to that described in Preparation 1.

30

¹H NMR (270MHz, CDCl₃) δ 7.36-7.29 (5H, m), 4.66 and 4.63, and 4.46 and 4.35 (total 2H, each d, J = 7.0Hz), 4.56 and 4.50(total 1H, each br.s), 4.27-4.20 (1H,

m), 3.81-3.14 (4H, m), 3.36 and 3.10 (total 3H, each s), 2.10-1.80 (2H, m), 2.00(2H, br.s).

IR (neat) : 3350, 3300, 1650 cm^{-1} .

Preparation 6

5 (2S,3S)-1-(2-Amino-2-phenylethyl)-3-(methoxymethoxy)pyrrolidine

This compound was prepared in 97.7% yield according to a procedure similar to that described in Preparation 2.

^1H NMR (270MHz, CDCl_3) δ 7.39-7.24 (5H, m), 4.66 (1H, d, $J = 6.6\text{Hz}$), 4.63 (1H, d, $J = 7.0\text{Hz}$), 4.30-4.23 (1H, m), 4.08 (1H, dd, $J = 3.3, 10.3\text{Hz}$), 3.37 (3H, s), 3.00-2.80 (2H, m), 2.74 (1H, dd $J = 10.3, 11.7\text{Hz}$), 2.60-2.45 (2H, m), 2.41 (1H, dd, $J = 3.3, 11.7\text{Hz}$), 2.21-2.04 (1H, m), 1.95-1.75 (3H, m).

IR (neat) : 3400 cm^{-1} .

Example 23

(2S,3S)-1-[2-N-(2,2-Difluoroethylamino)-2-phenylethyl]-3-hydroxypyrrolidine

15 To a stirred solution of (2S,3S)-1-(2-amino-2-phenylethyl)-3-(methoxymethoxy)pyrrolidine (0.77g, 3.08mmol) and 2,2-difluoroacetic acid (0.21ml, 3.3mmol) was added 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (0.63g, 3.3mmol) at room temperature. After 4h stirring, the reaction mixture was diluted with CH_2Cl_2 (20ml), washed with saturated NaHCO_3 aqueous solution and brine, dried (Na_2SO_4), and concentrated to give 1.05g of brown viscous oil. This was purified by column chromatography (silica gel:100g, $\text{CH}_2\text{Cl}_2/\text{MeOH} : 40/1$ as eluent) to give 0.69g (68.3%) of brown viscous oil.

25 ^1H NMR (270MHz, CDCl_3) δ 7.37-7.25 (5H, m), 5.91 (1H, t, $J_{\text{H,F}} = 54.2\text{Hz}$), 4.95-4.85 (1H, m), 4.63 (1H, d, $J = 6.6\text{Hz}$), 4.59 (1H, d, $J = 6.6\text{Hz}$), 4.26-4.19 (1H, m), 3.35 (3H, s), 2.96-2.69 (4H, m), 2.56-2.43 (2H, m), 2.18-2.03 (1H, m), 1.87-1.75 (1H, m), 1.72 (1H, br.s).

IR (neat) : 3300, 1700 cm^{-1} .

To a stirred solution of this amide derivative (0.69g, 2.1mmol) in THF (6ml) was added $\text{BH}_3 \cdot \text{Me}_2\text{S}$ (0.6ml, 6.3mmol) at room temperature and then refluxed for 30 5h. After cooling down to room temperature, 1N HCl (10ml) was added dropwise carefully and the resulting mixture was refluxed for 2h. After cooling down to room temperature, the reaction mixture was basified with 1N NaOH aqueous solution to pH

12 and extracted with CH_2Cl_2 (20ml x 3). The extract was dried (Na_2SO_4) and concentrated to give 566mg (67.9% for two steps) of titled compound as yellow viscous oil.

^1H NMR (270MHz, CDCl_3) δ 7.37-7.23 (5H, m), 5.76 (1H, ddt, $J = 3.3, 5.1, 56.1\text{Hz}$), 4.40-4.25 (1H, m), 3.80 (1H, dd, $J = 3.7, 10.6\text{Hz}$), 3.77-3.70 (1H, m), 2.99 (1H, dt, $J = 5.5, 8.8\text{Hz}$), 2.95-2.45 (4H, m), 2.82 (1H, dd, $J = 11.0, 12.1\text{Hz}$), 2.36-1.95 (3H, m), 1.90-1.70 (2H, m).
 IR (neat) : 3400, 3330 cm^{-1} .

Example 24

10 2-(3,4-Dichlorophenyl)-N-(2,2-difluoroethyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This compound was prepared in 62.7% yield according to a procedure similar to that described in Example 1.

^1H NMR (270MHz, CDCl_3) δ 7.43-7.26 (6H, m), 7.15-7.09 (2H, m), 6.13(0.5H, dd, $J = 5.5, 11.0\text{Hz}$), 6.03 (0.5H, ddt, $J = 2.9, 5.1, 55.0\text{Hz}$), 5.27 (0.5H, ddt, $J = 4.4, 4.8, 56.4\text{Hz}$), 5.05(0.5H, dd, $J = 6.6, 7.7\text{Hz}$), 4.35-4.25 (1H, m), 3.86 (1H, s), 3.84 (0.5H, d, $J = 15.8\text{Hz}$), 3.75(0.5H, d, $J = 15.8\text{Hz}$), 3.70-2.60 (7H, m), 1.80-1.65 (1H, m).

IR (neat) : 3450, 1650 cm^{-1} .

20 HCl salt : amorphous solid

Anal. Calcd for $\text{C}_{22}\text{H}_{24}\text{Cl}_2\text{F}_2\text{N}_2\text{O}_2 \cdot \text{HCl} \cdot \text{H}_2\text{O}$: C, 51.63 ; H, 5.32 ; N, 5.47.

Found : C, 51.94 ; H, 5.40 ; N, 5.51.

Preparation 7

25 (2S,3S)-1-[2-N-(2-Fluoroethylamino)-2-phenylethyl]-3-(methoxymethoxy)pyrrolidine

A mixture of (2S,3S)-1-(2-amino-2-phenylethyl)-3-(methoxymethoxy)pyrrolidine (1.01g, 4mmol), 2-bromofluoroethane (0.90ml, 12mmol), and K_2CO_3 (0.69g, 5mmol) in DMF (5ml) was stirred at 70 for 13h. The reaction mixture was diluted with water (10ml), basified with 1N NaOH aqueous solution to pH12, and extracted with CH_2Cl_2 (30ml x 3). The extract was dried (Na_2SO_4) and concentrated in vacuo to give 1.00g of dark brown viscous oil, which was purified by column chromatography (silica gel:100g, $\text{CH}_2\text{Cl}_2/\text{MeOH}$: 30/1 as

eluent) to afford 0.51g (43%) of brown viscous oil.

¹H NMR (270MHz, CDCl₃) δ 7.38-7.25 (5H, m), 4.66 (1H, d, J = 7.0Hz), 4.63 (1H, d, J = 7.0Hz), 4.60-4.22 (3H, m), 3.77 (1H, dd, J = 3.3, 10.3Hz), 3.38 (3H, s), 2.98 (1H, dd, J = 6.2, 10.3Hz), 2.94-2.49 (6H, m), 2.34 (1H, dd, J = 3.7, 12.1Hz), 2.20-2.06 (1H, m), 1.90-1.78 (1H, m), 1.56 (1H, br.s).
 IR (neat) : 3320cm⁻¹.

Example 25

2-(3,4-Dichlorophenyl)-N-(2-fluoroethyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

To a stirred solution of (2S,3S)-1-[2-N-(2-fluoroethylamino)-2-phenylethyl]-3-(methoxymethoxy)pyrrolidine (0.7g, 2.6mmol) and triethylamine (0.56ml, 4mmol) in CH₂Cl₂ (10ml) was added 3,4-dichlorophenylacetyl chloride [this was prepared from 3,4-dichlorophenylacetic acid (0.82g, 4mmol) and thionyl chloride (0.36ml, 5mmol)] at room temperature. After 20min, the reaction mixture was diluted with CH₂Cl₂ (20ml), washed with saturated NaHCO₃ aqueous solution. and dried (Na₂SO₄). Evaporation of the solvent gave 1.96g of brown oil, which was purified by column chromatography (silica gel 100g, CH₂Cl₂/MeOH : 20/1 as eluent) to give 0.81g (64.3%) of yellow viscous oil. To this oil was added HCl gas saturated MeOH solution (10ml) and stirred at room temperature for 2h. The solvent was evaporated and the residue was basified with 1N NaOH aqueous solution to pH 11 and extracted with CH₂Cl₂ (20ml x 2). After dry (Na₂SO₄), the solvent was evaporated to afford 0.78g of brown viscous oil, which was purified by column chromatography (silica gel : 100g, CH₂Cl₂/MeOH : 40/1 as eluent) to give 0.598g (81.5%) of clear yellow viscous oil.

¹H NMR (270MHz, CDCl₃) δ 7.41-7.26 (6H, m), 7.20-7.09 (2H, m), 6.10 (0.6H, dd, J = 5.9, 10.6Hz), 5.03 (0.4H, t, J = 7.3Hz), 4.70-3.90 (3H, m), 3.83 (0.6H, d, J = 15.4Hz), 3.83 (0.8H, s), 3.75 (0.6H, d, J = 15.8Hz), 3.72-3.25 (2H, m), 3.22-2.98 (2H, m), 2.94-2.80 (1H, m), 2.75 -2.08 (5H, m), 1.80-1.65 (1H, m).
 IR (neat) : 3400, 1640cm⁻¹.

HCl salt : amorphous solid.

Anal. Calcd for C₂₂H₂₅Cl₂FN₂O₂•HCl•2H₂O : C, 51.63; H, 5.91; N, 5.47.

Found : C, 51.95; H, 5.64; N, 5.42.

Example 26**(2S,3S)-1-(2-N-Benzylamino-2-phenylethyl)-3-hydroxypyrrolidine**

This compound was prepared in 100% yield according to a procedure similar to that described in Example 9.

- 5 ^1H NMR (270MHz, CDCl_3) δ 7.43-7.22 (10H, m), 4.30- 4.25 (1H, m), 3.77 (1H, d, $J = 13.6\text{Hz}$), 3.73 (1H, dd, $J = 3.7, 11.0\text{Hz}$), 3.49 (1H, d, $J = 13.6\text{Hz}$), 2.87 (1H, dd, $J = 11.0, 12.1\text{Hz}$), 2.85-2.75 (1H, m), 2.64-2.51 (2H, m), 2.38 (2H, br.s), 2.30 (1H, dd, $J = 3.7, 12.1\text{Hz}$), 2.23-2.07 (2H, m), 1.77-1.65 (1H, m).
 1R (neat) : 3300cm^{-1} .

10

Example 27**N-Benzyl-2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxy-pyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide**

This compound was prepared in 62.6% yield according to a procedure similar to that described in Example 1.

- 15 ^1H NMR (270MHz, CDCl_3) δ 7.40-6.78 (13H, m), 6.27 (1H, br.d, $J = 8.1\text{Hz}$), 4.93 (1H, br.d, $J = 17.6\text{Hz}$), 4.65-4.50 (2H, m), 4.20-3.95 (2H, m), 3.85-3.60 (3H, m), 3.55-3.15 (4H, m), 2.50-2.35 (1H, m), 2.30-2.10 (1H, m).
 1R (KBr) : $3330, 1640\text{cm}^{-1}$.

HCl salt : amorphous solid

- 20 Anal. Calcd for $\text{C}_{27}\text{H}_{28}\text{Cl}_2\text{N}_2\text{O}_2 \cdot \text{HCl} \cdot \text{H}_2\text{O}$: C, 60.29 ; H, 5.81 ; N, 5.21.
 Found : C, 60.49 ; H, 5.38 ; N, 5.24.

Example 28**N-Carbamoylmethyl-2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxy-pyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide**

- 25 A mixture of N-cyanomethyl-2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxy-pyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide(2.54g, 5.9mmol) and HCl gas saturated methanol(20ml) was stirred at room temperature for 0.5h. The reaction mixture was concentrated, basified with aqueous NH_3 solution, and extracted with CH_2Cl_2 (30ml x 3). After dry(Na_2SO_4), the solvent was evaporated to give brown viscous oil, which
 30 was purified by column chromatography(silica gel:100g, CH_2Cl_2 /MeOH: 30/1 to 15/1 as eluent) to afford 1.25g(47.3%) of pale yellow viscous oil.

^1H NMR (270MHz, CDCl_3) δ 10.14 (1H, br.s), 7.40-7.09 (8H, m), 6.31 (1H, dd,

$J = 3.3, 12.8\text{Hz}$), 5.70 (1H, br.s), 4.42 (1H, m), 3.80-3.55 (4H, m), 3.50-3.40 (1H, m), 3.35 (1H, app t, $J = 12.8\text{Hz}$), 2.76 (1H, $J = 10.3\text{Hz}$), 2.75-2.64 (2H, m, including 1H, dd, $J = 3.3, 12.8\text{Hz}$ at 2.67ppm), 2.30-2.15 (2H, m), 1.90-1.70 (2H, m).

5 ^{13}C NMR (CDCl_3) δ 173.2, 172.1, 137.4, 134.5, 132.5, 131.1, 130.4, 128.94, 128.86, 128.6, 128.3, 127.1, 70.4, 63.1, 53.9, 53.4, 51.4, 47.7, 39.8, 34.7.

1R (KBr) : 3400, 1690, 1650 cm^{-1} .

MS m/e (%) : 450 (3)[M+], 431 (2), 429 (3), 408 (2), 406 (3), 363 (4), 243 (3), 189 (75), 163 (99), 161 (94), 159 (99), 132 (36), 125 (55), 118 (100), 104 (95), 101 (98), 91 (91), 89 (70), 82 (61).

HCl salt : amorphous solid

Anal. Calcd for $\text{C}_{22}\text{H}_{25}\text{Cl}_2\text{N}_3\text{O}_3 \cdot \text{HCl} \cdot \text{H}_2\text{O}$: C, 52.34 ; H, 5.59 ; N, 8.32.

Found : C, 52.41 ; H, 5.38 ; N, 7.96.

15

Example 29

(2S,3S)-3-Hydroxy-1-[2-N-(2-methoxyethylamino)-2-phenylethyl]pyrrolidine

To a stirred solution of (2S,3S)-1-(2-amino-2-phenylethyl)-3-hydroxypyrrolidine (0.619g, 3mmol) and 2-methoxyacetic acid (0.23ml, 3mmol) in CH_2Cl_2 (10ml) was added 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (0.863g, 4.5mmol) at room temperature. After 0.5h stirring, the reaction mixture was diluted with water (50ml) and extracted with CH_2Cl_2 (30ml x 3). The extract was washed with saturated NaHCO_3 aqueous solution and brine, dried (Na_2SO_4), and concentrated to give 0.711g of clear yellow viscous oil. This was purified by column chromatography (silica gel: 40g, $\text{CH}_2\text{Cl}_2/\text{MeOH}$: 30/1 as eluent) to give 0.44g (52.7%) of pale yellow viscous oil.

25 ^1H NMR (270MHz, CDCl_3) δ 7.35-7.22 (6H, m), 5.07-4.99 (1H, m), 4.30-4.20 (1H, m), 3.91 (2H, s), 3.42 (3H, s), 3.02 (1H, s), 2.94-2.85 (2H, m), 2.67 (1H, dd, $J=5.1, 12.5\text{Hz}$), 2.60 (2H, d, $J=4.4\text{Hz}$), 2.36-2.27 (1H, m), 2.17-2.07 (1H, m), 1.71-1.66 (1H, m).

30 1R (neat) : 3350, 3270, 1660 cm^{-1} .

To a stirred suspension of lithium aluminum hydride (0.24g, 6mmol) in THF (10ml) was added a solution of this amide derivative (0.44g, 1.58mmol) in

THF(20ml) dropwise at room temperature and then the mixture was refluxed for 3.5h. After cooling down to room temperature, $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ (2.00g) and KF(0.2g) was added to the reaction mixture. After 20min stirring, the solid appeared was removed by Celite filtration. The filtrate was concentrated and the residue was purified by column chromatography (silica gel:12g, $\text{CH}_2\text{Cl}_2/\text{MeOH}$: 30/1 as eluent) to give 0.118g (28.2%) of pale yellow viscous oil.

^1H NMR (270MHz, CDCl_3) δ 7.39-7.23 (5H, m), 4.35-4.26 (1H, m), 3.78 (1H, dd, $J = 3.7, 10.6\text{Hz}$), 3.53-3.40 (2H, m), 3.35 (3H, s), 3.10-3.00 (1H, m), 2.95-2.85 (3H, m), 2.87-2.60 (4H, m), 2.38 (1H, dd, $J = 3.9, 12.1\text{Hz}$), 2.33-2.15 (2H, m), 1.85-1.73 (1H, m).

IR (neat) : 3400, 3330 cm^{-1} .

Example 30

2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(2-methoxyethyl)acetamide

This compound was prepared in 77.2% yield according to a procedure similar to that described in Example 1.

^1H NMR (270MHz, CDCl_3) δ 7.40-7.25 (6H, m), 7.18-7.05 (2H, m), 5.93 (0.8H, dd, $J = 5.9, 9.9\text{Hz}$), 5.00 (0.2H, t), 4.35-4.25 (1H, m), 3.82 (1H, d, $J = 15.4\text{Hz}$), 3.75 (1H, d, $J = 15.4\text{Hz}$), 3.55-2.55 (13H, m, including 2.4H, s, at 3.19ppm), 2.50-2.35 (1H, m), 2.25 -2.10 (1H, m), 1.85-1.70(1H, m).

IR (neat) : 3400, 1640 cm^{-1} .

HCl salt : amorphous solid.

Anal. Calcd for $\text{C}_{23}\text{H}_{28}\text{Cl}_2\text{N}_2\text{O}_3 \cdot \text{HCl} \cdot 2.2\text{H}_2\text{O}$: C, 52.37; H, 6.38; N, 5.31.

Found : C, 52.29; H, 6.40; N, 5.32.

Example 31

(2S,3S)-3-Hydroxy-1-[2-N-(2-methylthioethylamino)-2-phenylethyl]pyrrolidine

This compound was prepared in 50.5% yield according to a procedure similar to that described in Example 29.

^1H NMR (270MHz, CDCl_3) δ 7.38-7.24 (5H, m), 4.35-4.26 (1H, m), 3.75 (1H, dd, $J = 3.7, 10.6\text{Hz}$), 3.10-3.00 (1H, m), 2.85 (1H, dd, $J = 10.6, 12.1\text{Hz}$), 2.75-2.55 (5H, m), 2.41-2.15 (6H, m, including 1H, dd, $J = 4.0, 12.1\text{Hz}$ at 2.38ppm), 2.05 (3H, s), 1.85-1.70 (1H, m).

1R (neat) : 3300cm⁻¹.

Example 32

2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(2-methylthioethyl)acetamide

5 This compound was prepared in 64.8% yield according to a procedure similar to that described in Example 1.

¹H NMR (270MHz, CDCl₃) δ 7.45-7.25 (6H, m), 7.18-7.10 (2H, m), 6.06 (0.6H, dd, J = 5.9, 10.3Hz), 5.02 (0.4H, dd, J=6.2, 8.4Hz), 4.38-4.25 (1H, m), 3.83 (0.8H, s), 3.76 (0.6H, d, J=15.4Hz), 3.68 (0.6H, d, J = 15.4Hz), 3.50-1.65 (16H, m, including 1.2H, s, at 1.99ppm and 1.8H, s, at 1.96ppm), 2.50-2.35 (1H, m), 2.25-2.10 (1H, m), 1.85-1.70(1H, m).

1R (neat) : 3450, 1640cm⁻¹.

HCl salt : mp 195-197.5 .

Anal. Calcd for C₂₃H₂₈Cl₂N₂O₂S HCl 0.5H₂O : C, 53.86; H, 5.90; N, 5.46.

15 Found : C, 54.08; H, 5.91; N, 5.39.

Example 33

(2S,3S)-1-[2-N-(2-N,N-Dimethylaminoethylamino)-2-phenylethyl]-3-metoxymetoxypyrrolidine

20 This compound was prepared in 53.1% yield according to a procedure similar to that described in Example 29.

¹H NMR (270MHz, CDCl₃) δ 7.39-7.20 (5H, m), 4.65 (1H, d, J=7.0Hz), 4.62 (1H, d, J=6.6Hz), 4.28-4.21 (1H, m), 3.70 (1H, dd, J = 3.7, 10.6Hz), 3.37 (3H, s), 2.95 (1H, dd, J=6.2, 9.9Hz), 2.82 (1H, dd, J=10.6, 11.7Hz), 2.80-2.71 (1H, m), 2.59-2.45 (4H, m), 2.42-2.30 (4H, m), 2.18 (6H, s), 2.16-2.05 (1H, m), 1.85-1.75 (1H, m).

25 1R (neat) : 3300cm⁻¹.

Example 34

2-(3,4-Dichlorophenyl)-N-(2-N,N-dimethylaminoethyl-amino)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

30 This compound was prepared in 88% yield according to a procedure similar to that described in Examples 1 and 25.

¹H NMR (270MHz, CDCl₃) δ 7.41-7.08 (8H, m), 6.03 (0.7H, dd, J = 5.9,

9.9Hz), 5.01 (0.3H, dd, J=7.0, 7.3Hz), 4.32-4.22 (1H, m), 3.79 (0.6H, s), 3.77 (0.7H, d, J=15.4Hz), 3.70 (0.7H, d, J = 15.4Hz), 3.36 (0.6H, app.t, J=7.0Hz), 3.20 (1.4H, app.t, J=7.3Hz), 3.15-3.00 (2H, m), 2.95-2.80 (1H, m), 2.75 -2.50 (2H, m), 2.50-1.61 (6H, m), 2.11 (1.8H, s), 2.09 (4.2H, s).

5 IR (neat) : 3400, 1640cm⁻¹.

HCl salt : amorphous solid.

Anal. Calcd for C₂₄H₃₁Cl₂N₃O₂ 2HCl 1.3H₂O : C, 51.40; H, 6.40; N, 7.49.

Found : C, 51.79; H, 7.01; N, 7.58.

Example 35

10 (2S,3S)-1-[2-N-(2,2-Dimethoxyethylamino)-2-phenylethyl]-3-hydroxypyrrolidine

This compound was prepared in 66% yield according to a procedure similar to that described in Example 2.

¹H NMR (270MHz, CDCl₃) δ 7.38-7.22 (5H, m), 4.45 (1H, t, J=5.1Hz), 4.33-4.27 (1H, m), 3.74 (1H, dd, J = 3.7, 10.6Hz), 3.36 (3H, s), 3.30 (3H, s), 3.08-3.00 (1H, m), 2.85 (1H, dd, J=10.6, 12.1Hz), 2.71 (1H, br.d, J=9.5Hz), 2.65-2.55 (3H, m, including 2H, d, J=5.1Hz at 2.61ppm), 2.37 (1H, dd, J=3.7, 12.1Hz), 2.32-2.12 (4H, m), 1.81-1.72 (1H, m).

15 IR (neat) : 3400, 3300cm⁻¹.

Example 36

20 2-(3,4-Dichlorophenyl)-N-(2,2-dimethoxyethyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This compound was prepared in 91.2% yield according to a procedure similar to that described in Example 29.

¹H NMR (270MHz, CDCl₃) δ 7.41-7.10 (8H, m), 6.09 (1H, dd, J = 5.9, 9.9Hz), 4.35-4.25 (1H, m), 3.94 (1H, d, J=15.4Hz), 3.85 (1H, d, J=15.8Hz), 3.59 (1H, t, J=5.1Hz), 3.34-3.15 (3H, m), 3.22 (3H, s), 3.18 (3H, s), 3.10-3.00 (1H, m), 2.86 (1H, dd, J=5.9, 12.5Hz), 2.76-2.63 (2H, m), 2.40-2.25 (1H, m), 2.20-2.08 (1H, m), 1.90-1.65 (2H, m).

25 IR (neat) : 3450, 1640cm⁻¹.

30 Fumalic acid salt : amorphous solid.

Anal. Calcd for C₂₄H₃₀Cl₂N₂O₄ C₄H₄O₄ 2H₂O : C, 53.09; H, 6.05; N, 4.92.

Found : C, 53.47; H, 6.04; N, 4.51.

Example 37**(2S,3S)-3-Hydroxy-1-[2-phenylethyl]-2-N-(2-pyrrolyl)methyl-aminolpyrrolidine**

This compound was prepared in 100% yield according to a procedure similar to that described in Example 9.

- 5 ^1H NMR (270MHz, CDCl_3) δ 9.29 (1H, br.s), 7.37-7.25 (5H, m), 6.75 (1H, d, $J=1.5\text{Hz}$), 6.12 (1H, dd, $J=2.6, 5.5\text{Hz}$), 5.97 (1H, br.s), 4.35-4.25 (1H, m), 3.78 (1H, d, $J=13.9\text{Hz}$), 3.73 (1H, dd, $J=3.3, 11.3\text{Hz}$), 3.55 (1H, d, $J=13.9\text{Hz}$), 3.31 (2H, br.s), 2.92 (1H, dd, $J=11.7, 12.1\text{Hz}$), 2.87-2.77 (2H, m), 2.52 (1H, dd, $J=4.8, 9.9\text{Hz}$), 2.30 (1H, dd, $J=3.3, 12.1\text{Hz}$), 2.25-2.02 (2H, m), 1.83-1.73 (1H, m).
- 10 IR (neat) : 3300cm^{-1} .

Example 38**2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(2-pyrrolyl)methylacetamide**

- 15 This compound was prepared in 81.3% yield according to a procedure similar to that described in Example 29.
- ^1H NMR (270MHz, CDCl_3) δ 12.00 (1H, br.s), 7.45-7.25 (6H, m), 7.08 (1H, d, $J=1.5\text{Hz}$), 6.87 (1H, dd, $J=1.5, 8.1\text{Hz}$), 6.80-6.75 (1H, m), 6.38 (1H, dd, $J=3.7, 12.8\text{Hz}$), 6.13 (1H, dd, $J=2.6, 5.1\text{Hz}$), 5.99 (1H, br.s), 4.58-4.50 (1H, m), 4.15 (2H, s), 3.65-3.40 (4H, m), 3.00-2.90 (2H, m), 2.80-2.65 (2H, m), 2.40-2.25 (2H, m), 1.90-1.65 (2H, m).
- 20 IR (neat) : 3450, 1630cm^{-1} .

Fumalic acid salt : amorphous solid.

Anal. Calcd for $\text{C}_{25}\text{H}_{27}\text{Cl}_2\text{N}_3\text{O}_2 \cdot \text{C}_4\text{H}_4\text{O}_4 \cdot \text{H}_2\text{O}$: C, 57.43; H, 5.48; N, 6.93.

- 25 Found : C, 57.46; H, 5.41; N, 6.95.

Example 39**(2S,3S)-3-Hydroxy-1-[2-N-(1-methyl-2-pyrrolyl)methylamino-2-phenylethyl]pyrrolidine**

- This compound was prepared in 100% yield according to a procedure similar to that described in Example 9.
- 30

^1H NMR (270MHz, CDCl_3) δ 7.43-7.24 (5H, m), 6.55 (1H, dd, $J=1.8, 2.6\text{Hz}$), 6.05-5.99 (2H, m), 4.30-4.25 (1H, m), 3.75 (1H, dd, $J=3.7, 10.6\text{Hz}$), 3.61 (1H, d,

$J=13.9\text{Hz}$), 3.55 (3H, s), 3.49 (1H, d, $J=13.6\text{Hz}$), 2.87-2.81 (1H, m), 2.81 (1H, dd, $J=10.6, 12.1\text{Hz}$), 2.62-2.53 (2H, m), 2.29 (1H, dd, $J=3.7, 12.1\text{Hz}$), 2.27-2.08 (2H, m), 1.99 (2H, br.s), 1.76-1.65 (1H, m).

IR (neat) : 3250cm^{-1} .

5

Example 40

2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(1-methyl-2-pyrrolyl)methyl-acetamide

This compound was prepared in 57.6% yield according to a procedure similar to that described in Example 29.

10 ^1H NMR (270MHz, DMSO- d_6) δ 7.54 (1H, d, $J=8.1\text{Hz}$), 7.39 (1H, br.s), 7.35-7.20 (5H, m), 7.17 (1H, br.d, $J=8.4\text{Hz}$), 6.59 (1H, br.s), 5.85-5.70 (2H, m), 5.60-5.50 (1H, m), 4.75-4.60 (1H, m), 4.42 (2H, ABq), 4.13 (1H, br.s), 3.70 (1H, s), 3.41 (3H, s), 3.10-2.75 (3H, m), 2.65-2.20 (4H, m), 2.00-1.85 (1H, m), 1.55-1.45 (1H, m).

15 IR (neat) : 3450, 1640cm^{-1} .

Fumalic acid salt : amorphous solid.

Anal. Calcd for $\text{C}_{25}\text{H}_{27}\text{Cl}_2\text{N}_3\text{O}_2 \cdot \text{C}_4\text{H}_4\text{O}_4 \cdot 1.5\text{H}_2\text{O}$: C, 57.24; H, 5.76; N, 6.67.

Found : C, 57.17; H, 5.40; N, 6.52.

Example 41

20 (2S,3S)-3-Hydroxy-1-[2-N-(methoxycarbonyl)methylamino-2-phenylethyl]pyrrolidine

This compound was prepared in 81.4% yield according to a procedure similar to that described in Preparation 3.

25 ^1H NMR (270MHz, CDCl_3) δ 7.36-7.24 (5H, m), 4.37-4.30 (1H, m), 3.82 (1H, dd, $J=3.7, 11.0\text{Hz}$), 3.69 (3H, s), 3.39 (1H, d, $J=17.9\text{Hz}$), 3.22-3.15 (1H, m), 3.18 (1H, d, $J=17.6\text{Hz}$), 2.86 (1H, dd, $J=11.0, 12.1\text{Hz}$), 2.73 (1H, dd, 4.8, 9.9Hz), 2.66 (1H, br.d, $J=8.1\text{Hz}$), 2.60-2.15 (5H, m, including 1H, dd, $J=3.7, 12.1\text{Hz}$, at 2.32ppm), 1.82-1.75 (1H, m).

Example 42

30 2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-methoxycarbonylmethylacetamide

This compound was prepared in 57.6% yield according to a procedure similar

to that described in Example 29.

^1H NMR (270MHz, CDCl_3) δ 7.41-7.20 (6H, m), 7.16-7.08 (2H, m), 6.03 (0.5H, dd, $J=5.9, 10.3\text{Hz}$), 4.99 (0.5H, dd, $J=6.2, 8.8\text{Hz}$), 4.28 (0.5H, d, $J=16.9\text{Hz}$), 4.30-4.20 (1H, m), 3.90 (1H, s), 3.87 (0.5H, d, $J=16.9\text{Hz}$), 3.77 (0.5H, d, $J=12.5\text{Hz}$), 3.75 (1H, s), 3.72 (0.5H, d, $J=12.5\text{Hz}$), 3.68 (1.5H, s), 3.63 (1.5H, s), 3.16-2.42 (5H, m), 2.27-1.65 (4H, m).

IR (neat) : 3450, 1750, 1650 cm^{-1} .

HCl salt : mp 169-172 .

Anal. Calcd for $\text{C}_{23}\text{H}_{26}\text{Cl}_2\text{N}_2\text{O}_4 \cdot \text{HCl} \cdot 0.5\text{H}_2\text{O}$: C, 54.08; H, 5.52; N, 5.48.

10 Found : C, 54.39; H, 5.49; N, 5.53.

Example 43

(2S,3S)-1-[2-N-(2-Cyanoethylamino)-2-phenylethyl]-3-hydroxypyrrolidine

A mixture of (2S,3S)-1-(2-amino-2-phenylethyl)-3-hydroxypyrrolidine (0.619g, 3mmol) and acrylonitrile (1ml, 15mmol) in ethanol (6ml) was stirred at room temperature for 13h. The solvent was evaporated to give 0.786mg (100%) of a yellow solid.

^1H NMR (270MHz, CDCl_3) δ 7.39-7.24 (5H, m), 4.37-4.32 (1H, m), 3.76 (1H, dd, $J=3.7, 11.0\text{Hz}$), 3.08-3.00 (1H, m), 2.90-2.72 (3H, m, including 1H, dd, $J=11.0, 12.1\text{Hz}$, at 2.82ppm), 2.69 (2H, d, $J=3.7\text{Hz}$), 2.53-2.14 (7H, m, including 1H, dd, $J=3.7, 12.1\text{Hz}$ at 2.34ppm), 1.83-1.73 (1H, m),.

IR (neat) : 3350, 3280, 2250 cm^{-1} .

Example 44

N-2-Cyanoethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

25 This compound was prepared in 93.3% yield according to a procedure similar to that described in Example 29.

^1H NMR (270MHz, CDCl_3) δ 7.42-7.06 (8H, m), 6.11 (0.3H, dd, $J=4.8, 11.4\text{Hz}$), 5.03 (0.7H, dd, $J=5.9, 9.2\text{Hz}$), 4.40-4.28 (1H, m), 3.85-3.40 (4H, m, including 2H, br.s at 3.77ppm), 3.30-2.60 (5H, m), 2.45-2.08 (4H, m), 1.90 (1H, br.s), 1.80-1.65 (1H, m).

IR (neat) : 3450, 2250, 1650 cm^{-1} .


Maleic acid salt : amorphous solid.

Anal. Calcd for $C_{23}H_{25}Cl_2N_3O_2 \cdot C_4H_4O_4 \cdot H_2O$: C, 55.87; H, 5.38; N, 7.24.

Found : C, 56.12; H, 5.35 N, 7.26.

CLAIMS

1. A compound of the following formula:



- 5 and its pharmaceutically acceptable salt, wherein
- R is hydrogen or hydroxy;
- Ar is phenyl or phenyl substituted with one to three substituents selected from halo, C₁₋₄ alkyl and C₁₋₄ alkoxy;
- 10 X is phenyl or heterocyclic; phenyl or heterocyclic substituted with one to three substituents selected from halo, C₁₋₄ alkyl, C₁₋₄ alkoxy and methoxycarbonyl; mono-, di- or tri-halomethyl; cyano; COR¹, CH=NOR², OR², SR², CH₂CN, CH₂OR², CH₂SR², CH₂S(O)R², CH₂S(O)₂R², CH₂(R²)R³, CH₂N(R²)R³, CH₂NR²OH, CH₂N(COR²)OH, CH₂NR²COR³, CH₂NR²S(O)₂R³ or CH₂OCOR², wherein R¹ is hydrogen,
- 15 hydroxy, amino, NHOH, NHOCH₃, pyridylamino, NHN(CH₃)₂, C₁₋₄ alkoxy, benzyloxy, C₁₋₄ alkylamino, di-C₁₋₄ alkylamino, C₁₋₄ alkyl or C₁₋₄ alkylthio; and R² and R³ are each hydrogen, C₁₋₄ alkyl, C₁₋₄ alkoxy or C₇₋₁₁ phenylalkyl; and
- 20 X¹ is phenyl, furyl, thienyl, pyridyl, thiazolyl, benzofuryl or benzothienyl; phenyl, furyl, thienyl, pyridyl, thiazolyl, benzofuryl or benzothienyl, substituted with one to three substituents selected from halo, C₁₋₄ alkyl, C₁₋₄ alkoxy, amino, hydroxy, nitro, trifluoromethyl and mesyl.
2. A compound according to Claim 1, wherein R is hydroxy; Ar is phenyl optionally substituted with one to three halogen atoms; and X¹ is phenyl optionally
- 25 substituted with one to three halogen atoms.
3. A compound according to Claim 2, wherein Ar is phenyl; X¹ is 3,4-dichlorophenyl.

4. A compound according to Claim 3, wherein X is phenyl optionally substituted with one to three substituents selected from halo, C₁₋₄ alkyl, C₁₋₄ alkoxy and methoxycarbonyl.

5. A compound according to Claim 3, wherein X is mono-, di- or tri-halomethyl, cyano, hydroxycarbonyl, butyloxycarbonyl, benzyloxycarbonyl, carbamoyl or hydroxymethyl.

6. A compound according to Claim 3, wherein X is heterocyclic selected from furyl, thienyl and pyridyl.

7. A compound according to Claim 1, being one of the following:

10 N-Carboxymethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;

2-(3,4-Dichlorophenyl)-N-(2-hydroxyethyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;

15 2-(3,4-Dichlorophenyl)-N-[(2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl)-N-(2,2,2-trifluoroethyl)acetamide;

2-(3,4-Dichlorophenyl)-N-furfuryl-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;

2-(3,4-Dichlorophenyl)-N-[2-(3-(s)-hydroxypyrrolidin-1-yl)-1-(s)-phenylethyl]-N-(4-pyridyl)methylacetamide;

20 2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(3-pyridyl)methylacetamide;

N-Cyanomethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;


25 2-(3,4-Dichlorophenyl)-N-(2,2-difluoroethyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;

N-2-cyanoethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide; and

2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-methoxycarbonylmethylacetamide.

30 8. A compound of the formula:

38



(II)

wherein

R is hydrogen or hydroxy;

Ar is phenyl or phenyl substituted with one to three substituents selected from halo,
 5 C₁₋₄ alkyl and C₁₋₄ alkoxy; and

X is phenyl or heterocyclic; phenyl or heterocyclic substituted with one to three
 substituents selected from halo, C₁₋₄ alkyl, C₁₋₄ alkoxy and methoxycarbonyl;
 mono-, di- or tri-halomethyl; cyano; COR¹, CH=NOR², OR², SR², CH₂CN,
 CH₂OR², CH₂SR², CH₂S(O)R², CH₂S(O)₂R², CH₂(R²)R³, CH₂N(R²)R³,
 10 CH₂NR²OH, CH₂N(COR²)OH, CH₂NR²COR³, CH₂NR²S(O)₂R³ or
 CH₂OCOR², wherein R¹ is hydrogen, hydroxy, amino, NHOH, NHOCH₃,
 pyridylamino, NHN(CH₃)₂, C₁₋₄ alkoxy, benzyloxy, C₁₋₄ alkylamino, di-C₁₋₄
 alkylamino, C₁₋₄ alkyl or C₁₋₄ alkylthio; and R² and R³ are each hydrogen,
 C₁₋₄ alkyl, C₁₋₄ alkoxy or C₇₋₁₁ phenylalkyl.

15 9. A compound according to Claim 8, wherein R is hydroxy; and Ar is
 phenyl.

10. A compound according to Claim 9, wherein X is phenyl,
 methoxycarbonylphenyl, mono-, di- or tri-halomethyl, cyano, hydroxycarbonyl,
 butyloxycarbonyl, benzyloxycarbonyl, carbamoyl, hydroxymethyl, furyl, thienyl or
 20 pyridyl.

11. A pharmaceutical composition useful as an analgesic, antiinflammatory,
 diuretic or neuroprotective agent, which comprises a compound according to claim
 1, and a pharmaceutically inert carrier.

INTERNATIONAL SEARCH REPORT

Intern al Application No
PCT/JP 94/01399A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D207/12 C07D207/32 C07D401/12 C07D405/12 C07D409/12
A61K31/40

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| Y | EP,A,0 483 580 (MERCK) 6 May 1992 see the whole document --- | 1-11 |
| Y | EP,A,0 569 802 (MERCK) 18 November 1993 see the whole document ----- | 1-11 |

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

6 April 1995

Date of mailing of the international search report

28. 04. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Kissler, B

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 94/01399

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date | |
|---|---------------------|----------------------------|---------------------|----------|
| EP-A-0483580 | 06-05-92 | DE-A- | 4034785 | 07-05-92 |
| | | AU-A- | 8553891 | 07-05-92 |
| | | CA-A- | 2054648 | 03-05-92 |
| | | JP-A- | 4264067 | 18-09-92 |
| | | US-A- | 5232978 | 03-08-93 |
| ----- | | | | |
| EP-A-0569802 | 18-11-93 | DE-A- | 4215213 | 11-11-93 |
| | | AU-B- | 3834193 | 11-11-93 |
| | | CN-A- | 1079219 | 08-12-93 |
| | | CZ-A- | 9300823 | 16-02-94 |
| | | JP-A- | 6049022 | 22-02-94 |
| | | PL-A- | 298845 | 10-01-94 |
| ----- | | | | |